

SEQUENCE LISTING

<110> ZHANG, HUANMIN

AX, ROY L

BELLIN, MARY E

<120> ISOLATED POLYNUCLEOTIDE SEQUENCES ENCODING A FERTILITY ASSOCIATED ANTIGEN

05051407101  
TOT 20445060  
<130> 210707US20

<150> US 60/218,140

<151> 2000-07-14

<160> 9

<170> PatentIn version 3.1

<210> 1

<211> 592

<212> DNA

<213> Bos sp.

<220>

<221> CDS

<222> (1)..(591)

<223>

[illegible]

Glu Ser Lys Ala Leu  
195

<210> 2

<211> 197

<212> PRT

<213> Bos sp.

<400> 2

Glu Lys Leu Asn Gly Asn Ser Arg Lys Gly Ile Thr Tyr Asn Tyr Val  
5 10 15

Le Ser Ser Arg Leu Gly Arg Asn Thr Tyr Lys Glu Gln Tyr Ala Phe  
20 25 30

Leu Tyr Lys Glu Lys Leu Val Ser Val Lys Gln Ser Tyr Leu Tyr His  
35 40 45

Asp Tyr Gln Ala Gly Asp Ala Asp Val Phe Ser Arg Glu Pro Phe Val  
50 55 60

Val Trp Phe Gln Ser Pro Tyr Thr Ala Val Lys Asp Phe Val Ile Val  
65 70 75 80

Pro Leu His Thr Thr Pro Glu Thr Ser Val Arg Glu Ile Asp Glu Leu  
85 90 95

Ala Asp Val Tyr Thr Asp Val Lys Arg Arg Trp Asn Ala Glu Asn Phe  
100 105 110

Ile Phe Met Gly Asp Phe Asn Ala Gly Cys Ser Tyr Val Pro Lys Lys  
115 120 125

Ala Trp Lys Asp Ile Arg Leu Arg Thr Asp Pro Lys Phe Val Trp Leu  
130 135 140

Ile Gly Asp Gln Glu Asp Thr Thr Val Lys Lys Ser Thr Asn Cys Ala  
 145 150 155 160

Tyr Asp Arg Ile Val Leu Arg Gly Gln Asn Ile Val Asn Ser Gly Gly  
 165 170 175

Pro Gln Ser Asn Leu Val Phe Asp Phe Gln Lys Ala Tyr Arg Leu Ser  
 180 185 190

Glu Ser Lys Ala Leu  
 195

<210> 3

<211> 671

<212> DNA

<213> Bos sp.

<240> 3

acacaggat ctgccccata ctgatggaga agctaaacgg aaattcaaga aaaggcataa 60  
 catacaacta tgtgattagc tctcgccttg gaagaaacac atataaagaa cagtatgcct 120  
 ttctctataa agaaaagcta gtgtctgtaa aacaaagcta cctctaccac gactatcagg 180  
 ctggagacgc agatgtgttt tccagggaac cctttgtggt ctggttccag tcaccctaca 240  
 ccgctgtcaa ggacttcgtg attgtcccc tgcacaccac ccctgagaca tccgttagag 300  
 agattgatga gctggctgat gtctacacag atgtgaaacg tcgctggaat gcagagaatt 360  
 tcattttcat gggtgacttc aatgctggct gcagctacgt cccaagaag gcctggaagg 420  
 acatccgcct gaggacggac cccaagtctg tttggctgat cggggaccaa gaggacacca 480  
 cggtaagaa gagcacaac tgcgcctatg acaggatcgt gcttagagga caaatattg 540  
 tcaactctgg tggctctcaa tcaaacctcg tctttgattt ccagaaagct tacaggttgt 600  
 ctgaatcgaa ggccctggat gtcagcgacc actttccagt tcatcatcat catcatcatg 660  
 aagaaccatg a 671

<210> 4  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> synthetic DNA

<400> 4  
cgtgaggagc ttcggcgaga g

21

<210> 5  
<211> 26  
<212> PRT  
<213> Bos sp.

<220>  
<221> misc\_feature  
<222> (4)..(4)  
<223> Xaa is any amino acid

<400> 5  
Leu Lys Ile Xaa Ser Phe Asn Val Arg Ser Phe Gly Glu Ser Lys Lys  
1 5 10 15

Ala Gly Phe Asn Ala Met Arg Val Ile Val  
20 25

<210> 6  
<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic DNA

<400> 6

acaacaggat ctgccccata ctgatg

26

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic DNA

<400> 7

caactggaa agtggtcgct gacat

25

<210> 8

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic DNA

<400> 8

acaacaggat ctgccccata ctgatgg

27

<210> 9

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic DNA

<400> 9

tcatggttct tcatgatgat gatgatgatg aactggaaag tggtcgctga catccag

57

09905114.071501



Figure 1. Depicted recombinant FAA (rFAA), produced from cloned partial cDNA of bovine FAA gene in *E. coli*, showing the comparative position of the segment corresponding to intact bovine FAA.






Figure 2. Percent increase in acrosome reaction for each treatment above the control level. FAA present represents a fertile bull with detectable FAA on sperm and FAA negative represents a non-fertile bull without detectable FAA on sperm. The fertile bull (FAA present) reacted better to heparin induction of capacitation/acrosome reactions. Addition of FAA (5 ug/mL) stimulated maximum increase of acrosome reactions for each bull.




Figure 3. Effects of heparin alone (10  $\mu\text{g}/\text{ml}$ ) or recombinant FAA (rFAA, 20  $\mu\text{g}/\text{ml}$ ) with heparin to stimulate acrosome reaction in washed sperm from four fertile bulls.




Figure 4. Dose-response comparisons ( $\mu\text{g/ml}$ ) of the 22kDa recombinant FAA (rFAA) added with heparin ( $10 \mu\text{g/ml}$ ) to washed sperm. Averages from two to five different ejaculates are presented as each datum point for each bull.

Thursday, July 06, 2000

1 ACAACAGGAT CTGCCCCATA CTGATGGAGA AGCTAAACGG AAATTCAAGA  
51 AAAGGCATAA CATACAACTA TGTGATTAGC TCTCGCCTTG GAAGAAACAC  
101 ATATAAAGAA CAGTATGCCT TTCTCTATAA AGAAAAGCTA GTGTCTGTAA  
151 AACAAAGCTA CCTCTACCAC GACTATCAGG CTGGAGACGC AGATGTGTTT  
201 TCCAGGGAAC CCTTTGTGGT CTGGTTCCAG TCACCCTACA CCGCTGTCAA  
251 GGACTTCGTG ATTGTCCCCC TGCACACCAC CCCTGAGACA TCCGTTAGAG  
301 AGATTGATGA GCTGGCTGAT GTCTACACAG ATGTGAAACG TCGCTGGAAT  
351 GCAGAGAATT TCATTTTCAT GGGTGACTTC AATGCTGGCT GCAGCTACGT  
401 CCCCAGAAG GCCTGGAAGG ACATCCGCCT GAGGACGGAC CCAAGTTCG  
451 TTTGGCTGAT CGGGGACCAA GAGGACACCA CGGTCAAGAA GAGCACAAAC  
501 TGCGCCTATG ACAGGATCGT GCTTAGAGGA CAAAATATTG TCAACTCTGG  
551 TGGTCCTCAA TCAAACCTCG TCTTTGATTT CCAGAAAGCT TACAGGTTGT  
601 CTGAATCGAA GGCCCTGGAT GTCAGCGACC ACTTTCCAGT TCATCATCAT  
651 CATCATCATG AAGAACCATG A

Notes: Upstream primer sequence;

**Codon sequence responsible for the rFAA product;**

Stop codon.

0990514.071601

**SECRET**

[illegible]